

ALGORITHM FOR GENERATING ORTHOGONAL MATRICES WITH RATIONAL ELEMENTS.

R. A. SHARIPOV

ABSTRACT. Special orthogonal $n \times n$ matrices with rational elements form the group $\text{SO}(n, \mathbb{Q})$, where $\mathbb{Q}$ is the field of rational numbers. A theorem describing the structure of an arbitrary matrix from this group is proved. This theorem yields an algorithm for generating such matrices by means of random number routines.

1. INTRODUCTION.

Orthogonal matrices from the orthogonal group $\text{O}(n, \mathbb{R})$ describe rotations and rotations with reflections in the n -dimensional Euclidean space $\mathbb{R}^n$. Pure rotations constitute another classical group $\text{SO}(n, \mathbb{R})$, which is a subgroup in $\text{O}(n, \mathbb{R})$. The following matrix represents an elementary rotation in the p - q coordinate plane:

$$O_{[q]}^{[p]}(\varphi) = \left\| \begin{array}{cccccc} \overbrace{1 \quad \dots \quad 0}^p & \dots & 0 & \dots & 0 & \dots & 0 \\ \vdots & \ddots & \vdots & & \vdots & & \vdots \\ 0 & \dots & \alpha & \dots & -\beta & \dots & 0 \\ \vdots & & \vdots & \ddots & \vdots & & \vdots \\ 0 & \dots & \beta & \dots & \alpha & \dots & 0 \\ \vdots & & \vdots & & \vdots & \ddots & \vdots \\ \underbrace{0 \quad \dots \quad 0}_{q-1} & \dots & 0 & \dots & 0 & \dots & 1 \end{array} \right\|. \quad (1.1)$$

Here $\alpha = \cos(\varphi)$, $\beta = \sin(\varphi)$, and φ is the rotation angle. Matrices of the form (1.1) can be treated as elements of $\text{SO}(2, \mathbb{R})$ embedded into $\text{SO}(n, \mathbb{R})$. The matrix

$$\Omega = \left\| \begin{array}{c} \boxed{O^*} \\ 0 \quad \dots \quad 0 \quad 1 \end{array} \right\| \quad (1.2)$$

with $O^* \in \text{SO}(n-1, \mathbb{R})$ then is an element of $\text{SO}(n-1, \mathbb{R})$ embedded into $\text{SO}(n, \mathbb{R})$. For orthogonal matrices over the field of reals $\mathbb{R}$ there is the following theorem.

2000 *Mathematics Subject Classification.* 11D09, 15-04, 15A33.

Theorem 1.1. *Each orthogonal matrix $O \in \text{SO}(n, \mathbb{R})$ can be represented as a product $O = O^{[1]}(\varphi_1) \cdot \dots \cdot O^{[n-1]}(\varphi_{n-1}) \cdot \Omega$, where the first $n-1$ terms are matrices of elementary rotations (1.1), and Ω is an orthogonal matrix of the form (1.2).*

The angles $\varphi_1, \dots, \varphi_{n-1}$ in the theorem 1.1 are known as Euler angles (see Chapter VII in [1] for Euler angles in the three-dimensional case $n = 3$). These angles are restricted by the inequalities

$$0 \leq \varphi_1 \leq 2\pi, \quad 0 \leq \varphi_i \leq \pi \quad \text{for } i = 2, \dots, n-1.$$

Applying the theorem 1.1 recursively to O , then to O^* in (1.2), and so on, we easily prove the following theorem.

Theorem 1.2. *Each orthogonal matrix $O \in \text{SO}(n, \mathbb{R})$ can be represented as a product of $n(n-1)/2$ matrices of elementary rotations (1.1).*

The theorem 1.1 is not valid for orthogonal matrices over the field of rational numbers. As for the theorem 1.2, I don't know if it is valid or not for $O \in \text{SO}(n, \mathbb{Q})$. However, there is an algorithm for constructing orthogonal matrices over the field of rational numbers. This algorithm is exhausting, i.e. each orthogonal matrix $O \in \text{SO}(n, \mathbb{Q})$ can be obtained by applying this algorithm.

2. STEREOGRAPHIC PROJECTION.

Let $\mathfrak{S}$ be the unit sphere in $\mathbb{R}^n$. We take the point S with the coordinates $(0, \dots, 0, -1)$ for the south pole on this sphere. Then the equatorial hyperplane α is given by the equation $x^n = 0$, where x^n is the n -th coordinate of a point of




Fig. 2.1

$\mathbb{R}^n$ (we use upper indices for coordinates of vectors and points according to Einstein's tensorial notation, which is popular in differential geometry and in general relativity). Let X be an arbitrary point of the unit sphere $\mathfrak{S}$ and let $[SX)$ be the ray starting at the south pole S and passing through the point $X \in \mathfrak{S}$. This ray crosses the equatorial hyperplane at some definite point Y (as shown on figure 2.1 for three dimensional case). For each X point Y is unique. If we denote by $\mathfrak{S}^\circ$ the unit sphere $\mathfrak{S}$ with the pinned off south pole S , then we get the mapping $f: \mathfrak{S}^\circ \rightarrow \alpha$ that maps $\mathfrak{S}^\circ$ onto equatorial hyperplane α .

This map is called the *stereographic projection*. It is bijective and smooth. It's very important for us that the stereographic projection is given by rational functions. Indeed, if $x^1, \dots, x^n$ are the coordinates of some point $X \in \mathfrak{S}^\circ$ and if $y^1, \dots, y^{n-1}$ are the coordinates of the point $Y = f(X)$, then for $y^1, \dots, y^{n-1}$, we get the following formulas:

$$y^s = \frac{x^s}{1 + x^n} \quad \text{for } s = 1, \dots, n-1. \quad (2.1)$$

The inverse mapping $f^{-1}: Y \rightarrow X$ is also given by rational functions. Indeed, for the n -th coordinate x^n we have the formula

$$x^n = \left(1 - \sum_{i=1}^{n-1} (y^i)^2\right) / \left(1 + \sum_{i=1}^{n-1} (y^i)^2\right). \quad (2.2)$$

For the other coordinates of the point X from (2.1) and (2.2) we derive

$$x^s = \frac{2y^s}{\left(1 + \sum_{i=1}^{n-1} (y^i)^2\right)}, \text{ where } s = 1, \dots, n-1. \quad (2.3)$$

Rationality of the expressions (2.1), (2.2), and (2.3) mean that rational points of the pinned sphere $\mathfrak{S}^\circ$ are in one-to-one correspondence with rational points of the equatorial hyperplane α . As for the south pole S , it is usually associated with the infinite point $Y = \infty$ on α . Indeed, passing to the limit $Y \rightarrow \infty$ in the formulas (2.2) and (2.3), one can get the coordinates of the point S .

3. ORTHOGONAL MATRICES AND ONB'S.

Let $O \in \text{SO}(n, \mathbb{Q})$ be some orthogonal matrix. Its columns can be treated as vectors in $\mathbb{Q}^n$. Let's denote them $\mathbf{e}_1, \dots, \mathbf{e}_n$. Orthogonality of O means that the transposed matrix O^T coincides with the inverse matrix O^{-1} :

$$O \cdot O^T = 1. \quad (3.1)$$

Written in explicit form the equality (3.1) means that $\mathbf{e}_1, \dots, \mathbf{e}_n$ are vectors of the unit length perpendicular to each other. They form a so called ONB (orthonormal basis) in $\mathbb{Q}^n$ with respect to the standard scalar product

$$(\mathbf{X}, \mathbf{Y}) = \sum_{i=1}^n x^i y^i.$$

Matrices from the special orthogonal group $\text{SO}(n, \mathbb{Q})$ obey the additional restriction

$$\det Q = 1. \quad (3.2)$$

For the basis vectors $\mathbf{e}_1, \dots, \mathbf{e}_n$ the restriction (3.2) means that $\mathbf{e}_1, \dots, \mathbf{e}_n$ form a *right oriented* ONB (or a *right* ONB). In three dimensional space $\mathbb{Q}^3$ this property can be visualized.

Orientation rule. Vectors $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ form a *right triple* in $\mathbb{Q}^3$ if when looking from the end point of the third vector $\mathbf{e}_3$ the shortest rotation from the first vector $\mathbf{e}_1$ to the second vector $\mathbf{e}_2$ is observed as a counterclockwise rotation.

In physics (in electricity and magnetism) this rule is formulated as the rule of a right screw and also as the rule of a left hand (see [2] and [3]). In multidimensional spaces $n > 3$ one cannot visualize the concept of left and right since a human has no visual experience of living in such spaces. However, one can understand it mathematically by means of the theory of determinants and skew-symmetric

polylinear forms. Indeed, one should first prescribe right orientation to the standard basis in $\mathbb{Q}^n$ composed by the vectors

$$\mathbf{E}_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \quad \mathbf{E}_2 = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{pmatrix}, \quad \dots, \quad \mathbf{E}_n = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix}, \quad (3.3)$$

then one should consider the transition matrix S relating the standard basis (3.3) with another basis $\mathbf{e}_1, \dots, \mathbf{e}_n$. Its components are defined by the equality

$$\mathbf{e}_i = \sum_{j=1}^n S_i^j \cdot \mathbf{E}_j, \quad i = 1, \dots, n. \quad (3.4)$$

Definition 3.1. A basis $\mathbf{e}_1, \dots, \mathbf{e}_n$ in $\mathbb{Q}^n$ is called *right oriented* if $\det S > 0$. Otherwise, if $\det S < 0$, this base is called *left oriented*.

Another method of defining orientation in $\mathbb{Q}^n$ uses skew-symmetric polylinear forms. Remember that a polylinear n -form in $\mathbb{Q}^n$ is a $\mathbb{Q}$ -numeric function with n vectorial arguments $\omega = \omega(\mathbf{X}_1, \dots, \mathbf{X}_n)$ which is linear with respect to each its argument. A form ω is called completely skew-symmetric if its value changes the sign under permutation of any pair of arguments in it. It is known that a completely skew-symmetric n -form in $\mathbb{Q}^n$ is determined uniquely up to a scalar factor. Therefore there is exactly one n -form ω normalized by the condition $\omega(\mathbf{E}_1, \dots, \mathbf{E}_n) = 1$. It is called the volume form. For the basis $\mathbf{e}_1, \dots, \mathbf{e}_n$ we have the equality

$$\omega(\mathbf{e}_1, \dots, \mathbf{e}_n) = \det S \cdot \omega(\mathbf{E}_1, \dots, \mathbf{E}_n),$$

where the matrix S is determined by the formula (3.4). This means that the value of the volume form ω can be used as a measure of orientation for bases in $\mathbb{Q}^n$.

Note that if a basis $\mathbf{e}_1, \dots, \mathbf{e}_n$ is composed by the columns of an orthogonal matrix O , the transition matrix S in (3.4) coincides with O . This means that constructing an orthogonal matrix $O \in \text{SO}(n, \mathbb{Q})$ is equivalent to choosing some right oriented ONB in $\mathbb{Q}^n$. One vector in this base (say last vector $\mathbf{e}_n$) can be constructed by means of the stereographic projection as described in section 2 above. This is the first step in our algorithm for constructing orthogonal matrices. Then we should complement it with other $n-1$ vectors which should be unit vectors perpendicular to each other and perpendicular to the vector $\mathbf{e}_n$ as well. Below we use the Cayley transformation for this purpose.

4. CAYLEY TRANSFORMATION.

Let A be a skew-symmetric $n \times n$ square matrix. It is known that all eigenvalues of a skew-symmetric matrix are purely imaginary numbers (some of them can be equal to zero, but they cannot be nonzero real numbers). Therefore $\det(1 - A)$ is nonzero. Let's consider the matrix $O = (1 + A) \cdot (1 - A)^{-1}$. Matrices $1 + A$ and $1 - A$ commute with each other, hence $(1 + A) \cdot (1 - A)^{-1} = (1 - A)^{-1} \cdot (1 + A)$. For this reason we write $O = (1 + A) \cdot (1 - A)^{-1}$ as a fraction

$$O = \frac{1 + A}{1 - A}. \quad (4.1)$$

The formula (4.1) is known as the Cayley transformation (see [4]). If A is skew-symmetric, as it is in our case, then O is orthogonal matrix with $\det O = 1$. The Cayley transformation defines a mapping $\text{so}(n, \mathbb{Q}) \rightarrow \text{SO}(n, \mathbb{Q})$ similar to the exponential mapping $\exp: \text{so}(n, \mathbb{R}) \rightarrow \text{SO}(n, \mathbb{R})$. But, in contrast to the exponential mapping, it is rational, which is worth for our purposes. The Cayley transformation is an injective mapping. Indeed, if a matrix O is obtained by means of the formula (4.1), one can recover the matrix A by means of the formula

$$A = \frac{O - 1}{O + 1}. \quad (4.2)$$

However, one cannot apply the formula (4.2) to an arbitrary matrix $O \in \text{SO}(n, \mathbb{Q})$. Matrices with eigenvalues $\lambda = -1$ are not suitable. This means that the Cayley transformation is not surjective. Below we modify the Cayley transformation and convert it into an algorithm able to produce each matrix $O \in \text{SO}(n, \mathbb{Q})$.

We define some special skew-symmetric matrix. Taking $n - 1$ rational numbers $y^1, \dots, y^{n-1}$, we denote by $A[y^1, \dots, y^{n-1}]$ the matrix of the form

$$A[y^1, \dots, y^{n-1}] = \begin{pmatrix} 0 & \dots & 0 & y^1 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \dots & 0 & y^{n-1} \\ -y^1 & \dots & -y^{n-1} & 0 \end{pmatrix}. \quad (4.3)$$

Substituting (4.3) into (4.1), we obtain the orthogonal matrix

$$O[y^1, \dots, y^{n-1}] = \frac{1 + A[y^1, \dots, y^{n-1}]}{1 - A[y^1, \dots, y^{n-1}]}. \quad (4.4)$$

By direct calculation one can find that the n -th column in the matrix (4.4) coincides with the unit vector $\mathbf{e}_n$ constructed by means of the stereographic projection in section 2. Its components are given by the formulas (2.2) and (2.3). Let's denote by $\mathbf{e}_1, \dots, \mathbf{e}_{n-1}$ other $n - 1$ columns of the matrix $O[y^1, \dots, y^{n-1}]$. Completed by the vector $\mathbf{e}_n$, they form an ONB in $\mathbb{Q}^n$. Here are the formulas for the components of the k -th vector $\mathbf{e}_k$, where $k \neq n$:

$$x^s = \frac{-2 y^k y^s}{\left(1 + \sum_{i=1}^{n-1} (y^i)^2\right)} \quad \text{for } s \neq k, s \neq n, \quad (4.5)$$

$$x^k = 1 - \frac{2 (y^k)^2}{\left(1 + \sum_{i=1}^{n-1} (y^i)^2\right)}, \quad x^n = \frac{-2 y^k}{\left(1 + \sum_{i=1}^{n-1} (y^i)^2\right)}.$$

We can pass to the limit $Y \rightarrow \infty$ in (2.2) and (2.3). However, we cannot pass to

this limit in (4.5). For the infinity point $Y = \infty$ we set by definition

$$O[\infty] = \begin{bmatrix} 1 & \dots & 0 & 0 & 0 \\ \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & \dots & 1 & 0 & 0 \\ 0 & \dots & 0 & -1 & 0 \\ 0 & \dots & 0 & 0 & -1 \end{bmatrix} \quad (4.6)$$

Theorem 4.1. *Each unit vector $\mathbf{e} \in \mathbb{Q}^n$ given by its stereographic coordinates $y^1, \dots, y^{n-1}$ is canonically associated with some orthogonal matrix $O \in \text{SO}(n, \mathbb{Q})$.*

The formulas (2.2), (2.3), (4.5), and (4.6) give an explicit proof of the theorem 4.1. By construction the vector $\mathbf{e}$ is the n -th column in the matrix $O = O[y^1, \dots, y^{n-1}]$.

Now let O be some arbitrary matrix from the special orthogonal group $\text{SO}(n, \mathbb{Q})$. We denote by $\mathbf{e}$ its n -th column. This is the unit vector with rational components $x^1, \dots, x^n$. Suppose that $y^1, \dots, y^{n-1}$ are its stereographic coordinates (see formula (2.1) in section 2). They are also rational numbers. Therefore we can construct the orthogonal matrix $O[y^1, \dots, y^{n-1}]$ as described above. As a result we get two orthogonal matrices O and $O[y^1, \dots, y^{n-1}]$ with the same n -th column in them. This is possible if and only if these matrices are bound by the relationship

$$O = O[y^1, \dots, y^{n-1}] \cdot \Omega, \quad (4.7)$$

where Ω is an orthogonal matrix of the form (1.2). Thus we have proved a theorem.

Theorem 4.2. *Each orthogonal matrix $O \in \text{SO}(n, \mathbb{Q})$ can be represented as a product (4.7), where Ω is a blockwise diagonal matrix determined by some element of the orthogonal group $\text{SO}(n-1, \mathbb{Q})$.*

The theorem 4.2 is an analog of the theorem 1.1, while the rational parameters $y^1, \dots, y^{n-1}$ are analogs of Euler angles $\varphi_1, \dots, \varphi_{n-1}$. Applying this theorem recursively, for $O \in \text{SO}(n, \mathbb{Q})$ we get the equality

$$O = O[y_1^1, \dots, y_1^{n-1}] \cdot O[y_2^1, \dots, y_2^{n-2}] \cdot \dots \cdot O[y_{n-1}^1]. \quad (4.8)$$

Theorem 4.3. *Each orthogonal matrix $O \in \text{SO}(n, \mathbb{Q})$ can be constructed by means of the algorithm described above in the form of a product (4.8).*

The theorem 4.3 is an analog of the theorem 1.2. Note that the number of rational parameters y_j^i in (4.8) is equal to $n(n-1)/2$. This is exactly the same number as in the statement of the theorem 1.2.

5. CONCLUDING REMARKS.

I am not a specialist in algebra and I am not a specialist in number theory. I have encountered the problem of generating orthogonal matrices with rational elements in designing computer programs for testing students (see [5]). Therefore it's quite possible that all of the above results are not new. However, I hope that being gathered in one paper and formulated as a computational algorithm they could be useful for practical purposes. In addition, I have collected some references (see [6–12]) related to the subject of the present paper.

10. ACKNOWLEDGEMENTS.

I am grateful to V. A. Yuryev for helpful discussions. I am also grateful to Russian Fund for Basic Research and to the Academy of Sciences of the Republic Bashkortostan for financial support in 2001.

REFERENCES

1. Kostrikin A. I., *Introduction to algebra*, Nauka publishers, Moscow, 1977.
2. Borovoy A. A., Finkelstein E. B., Heruvimov A. N., *Laws of electromagnetism*, Nauka publishers, Moscow, 1970.
3. Landsberg G. S. (ed.), *Elementary physics manual, Vol. 2*, Nauka publishers, Moscow, 1975.
4. Postnikov M. M., *Lie Groups and Lie algebras. Lectures on geometry, 5-th term*, Nauka publishers, Moscow, 1982.
5. Sharipov R. A., *Orthogonal matrices with rational components in composing tests for High School students*, Paper math.GM/0006230 in Electronic Archive at LANL¹ (2000).
6. Grave D. A., *Treatise on algebraic analysis*, vol. 1 and 2, Kiev, 1938–1939.
7. Smirnov G. P., *On the representation of zero by quadratic forms*, Transactions of Bashkir State University, vol. 20, issue 2, Ufa, 1965.
8. Smirnov G. P., *On the solution of some Diophantine equations containing quadratic forms*, Transactions of Bashkir State University, vol. 20, issue 1, Ufa, 1965.
9. Smirnov G. P., *Entire orthogonal matrices and methods of their construction*, Transactions of Bashkir State University, vol. 31, issue 3, Ufa, 1968.
10. Bruening J. T., Lohmeier T. R., Sebaugh Ch. L., *Symmetric Pythagorean triple preserving matrices*, Missouri Journal of Mathematical Sciences **13** (2001), no. 1.
11. Wojtowicz M., *Algebraic structures of some sets of Pythagorean triples*, Missouri Journal of Mathematical Sciences **13** (2001), no. 1.
12. Crasmareanu M., *A new method to obtain Pythagorean triple preserving matrices*, Missouri Journal of Mathematical Sciences **14** (2002), no. 3.

5 RABOCHAYA STREET, 450003 UFA, RUSSIA

CELL PHONE: +7(917)476 93 48

E-mail address: r-sharipov@mail.ru

R_Sharipov@ic.bashedu.ru

URL: <http://www.freetextbooks.narod.ru>

<http://sovlit2.narod.ru>

¹Electronic Archive at Los Alamos National Laboratory of USA (LANL). The archive is accessible through the Internet <http://arXiv.org>, it has the mirror site <http://ru.arXiv.org> at the Institute for Theoretical and Experimental Physics (ITEP, Moscow).